

Optical phonon induced frictional drag in coupled two-dimensional electron gases

Ben Yu-Kuang Hu

*Mikroelektronik Centret, Bygning 345Ø, Danmarks Tekniske Universitet, DK-2800 Lyngby,
Denmark*

(February 1, 2008)

Abstract

The role of optical-phonons in frictional drag between two adjacent but electrically isolated two-dimensional electron gases is investigated. Since the optical-phonons in III-V materials have a considerably larger coupling to electrons than acoustic phonons (which are the dominant drag mechanism at low T and large separations) it might be expected that the optical phonons will give a large effect high temperatures. The two key differences between optical and acoustic phonon mediated drag are: (1) the optical phonon mediated interlayer interaction is short-ranged due to their negligible group velocity at the Brillouin zone center; and (2) the typical momentum transfer for an optical phonon mediated scattering is relatively large. These considerations make optical phonon mediated drag difficult to see in single subband GaAs systems, but it may be possible to see the effect in double subband GaAs systems or single subband quantum wells in a material with a lower effective mass and lower optical phonon energy such as InSb.

I. INTRODUCTION

The study of the effects of interactions between electrons makes up a considerable part of current research effort in condensed matter physics. One system which is particularly well-suited system for the study of phenomena associated with electron-electron interactions is the two-dimensional electron gas (2DEG) in a semiconductor quantum well. By changing various parameters in the system, either during the fabrication process or by the appropriate application of gate biases, magnetic field, temperature, etc. during the experiment, the physical conditions of the system can be changed over a wide range of parameter space. Probing electrical, optical, thermal or other properties of these 2DEGs often reveals interesting effects due to interactions.

Recently, it has been possible to make a direct study of interparticle interactions in 2DEGs, based on an proposal made some time ago.^{1,2} Experimentalists have been able to fabricate electrically isolated quantum wells which are only several hundred Å's apart. Thus, the electrons in the adjacent wells are coupled (through Coulomb and phonon mediated forces), but are unable to tunnel from one side to the other. These remarkable structures allow the experimentalist to probe directly the interlayer interactions by doing a "simple" transport measurement; i.e., turning on a current in one layer, and measuring the voltage built up in the second.³⁻⁸ These experiments provide a unique opportunity for a transport measurement to study effects of interparticle interaction.

The most obvious interlayer interaction is of course the Coulomb coupling, which gives relatively featureless dependences (at zero magnetic field) as a function of temperature, relative density, etc.⁹ Experiments, however, have shown that there are interesting pronounced features in drag which are associated with the collective modes in the system, indicating that under the right conditions drag is particularly sensitive to presence of these modes. So far, the drag effects from both acoustic phonons and plasmons have been observed^{3,5,8} and studied theoretically,¹⁰⁻¹⁵ but to the best of our knowledge no work on optical phonon mediated drag has been published.

Looking first at acoustic phonons in drag, it may seem surprising that they can contribute so strongly to the drag transresistivity, given their very weak coupling to the electron gas (relative to the Coulomb interaction). It turns out that strong phase-space effects at particular energy and momentum transfers cause the acoustic phonons to dominate in spite of the weak coupling.¹⁵ An obvious question arises: if the weak acoustic phonons give such a large contribution, can the optical phonons with a much larger coupling (witness the fact that scattering from optical phonons produces a much larger resistivity than scattering off their acoustic counterparts) give a correspondingly giant drag rate when they come in at higher temperatures?

The answer depends on the type of system one is studying. There are two significant differences between acoustic and optical phonons. The first is that, unlike their acoustic counterparts, the optical phonons around the Brillouin zone center (those which are relevant for drag) have a very small group velocity. Thus, the range of the optical phonons is much shorter than the acoustic phonons. In fact the longitudinal polar optical phonon mediated interaction has spatially exactly the same short-ranged form as the out-of-plane Coulomb interaction in two-dimensional structures; *i.e.*, it is proportional to $\exp(-qd)$ where $\hbar q$ is the electron momentum transferred and d is the center-to-center interlayer separation. Furthermore, every optical phonon mediated scattering event involves a transfer of the energy of an optical phonon. Since this energy is relatively large, conservation of energy dictates that the momentum transfer $\hbar q$ must also be large, which together with the $\exp(-qd)$ form of the interaction implies that the optical phonon mediated drag effect is usually small. However, as shown below, under certain experimental circumstances the effect may be relatively large and clearly observable. Provided the substantial experimental hurdles can be surmounted, it may be possible to perform a room temperature drag experiment.

The paper is organized as follows. In Section II, the formalism for optical phonon mediated drag in multiple subband systems is presented. In Section III the details of the calculation and the results are presented, both for the case of a single and double subband system. Finally, the conclusions are presented in Section IV.

II. FORMALISM

In this section, the generalization for the expression for the drag transresistivity for the case of several subbands is given. Then, the explicit form for polar optical phonon mediated interaction is introduced, following the formalism for phonon-mediated drag given by Zhang and Takahashi¹¹ and Bønsager et al.¹⁵

A. Transresistivity with multiple subband systems

In a typical drag experiment, two electrically isolated 2DEGs are placed close together. A current is passed through one layer (denoted layer 1), which induces a voltage in the second (layer 2). The quantity experimentalists measure is the transresistivity $\vec{\rho}_{21}$, which is the ratio of the electric field in the second layer to the current density in the first layer

$$\vec{\rho}_{21} \cdot \mathbf{J}_1 = \mathbf{E}_2. \quad (1)$$

Naturally, the stronger the interlayer coupling, the larger the magnitude of ρ_{21} .

We define the plane of the 2DEGs as the xy plane. We use upper case letters for 3-dimensional vectors and lower case for vectors in the plane of the 2DEG; e.g., $\mathbf{Q} = (\mathbf{q}, Q_z)$. The expression for the linear response transresistivity for multiple subbands systems, in the semi-classical weak interlayer interaction approximation, comes about from a simple generalization of the derivation in Ref. 14. At temperature T , with density n_i and charge e_i of layer i

$$\vec{\rho}_{21} = -\frac{\hbar^2}{2\pi e_1 e_2 n_1 n_2 k_B T} \int \frac{d\mathbf{q}}{(2\pi)^2} \int_0^\infty d\omega \sum_{\alpha\beta\gamma\delta} \frac{\mathbf{F}_1(\alpha\beta; \mathbf{q}, \omega) \mathbf{F}_2(\gamma\delta; \mathbf{q}, \omega) |v_{12}(\alpha\beta, \gamma\delta; \mathbf{q}, \omega)|^2}{\sinh^2(\hbar\omega/2k_B T)}. \quad (2)$$

Here $v_{12}(\alpha\beta\gamma\delta; \mathbf{q}, \omega)$ is the effective interlayer interaction which scatters carriers from subband $\alpha \rightarrow \beta$ in layer 1 and $\gamma \rightarrow \delta$ in layer 2, with the change of in-plane momentum $\hbar\mathbf{q}$ and energy $\hbar\omega$ (see Fig. 1). The functions $\mathbf{F}$ are given by

$$\mathbf{F}(\alpha\beta; q, \omega) = \frac{2\pi e}{\hbar\mu_t} \int \frac{d\mathbf{k}}{(2\pi)^2} [f_{0\alpha}(\mathbf{k}) - f_{0\beta}(\mathbf{k} + \mathbf{q})] \delta(\varepsilon_\alpha(\mathbf{k}) - \varepsilon_\beta(\mathbf{k} + \mathbf{q}) - \hbar\omega) \times \\ (\mathbf{v}_\alpha(\mathbf{k} + \mathbf{q})\tau_\alpha(\mathbf{k} + \mathbf{q}) - \mathbf{v}_\beta(\mathbf{k})\tau_\beta(\mathbf{k})) \quad (3)$$

where μ_t is the mobility of the sample, $\mathbf{v}_\alpha$ is the velocity, τ_α is the relaxation time and $f_{0\alpha}$ is the Fermi-Dirac distribution function in subband α . This relaxation time is defined by the linear response δf to a small electric field $\mathbf{E}$,

$$\delta f_\alpha(\mathbf{k}) = \left(-\frac{\partial f_{0\alpha}}{\partial \varepsilon} \right) e_i \mathbf{E} \cdot \mathbf{v}_\alpha(\mathbf{k}) \tau_\alpha(\mathbf{k}). \quad (4)$$

For parabolic isotropic systems where intralayer electron-electron scattering dominates over other scattering mechanisms, the distribution functions in all subbands are drifted Fermi-Diracs.^{14,16} Then, the expression for the transresistivity becomes

$$\rho_{21} = -\frac{h}{e^2} \frac{1}{4\pi} \int_0^\infty \frac{dq}{k_F^4} q^3 \int_0^\infty \frac{\hbar d\omega}{k_B T} \sum_{\alpha\beta\gamma\delta} \frac{\chi_1''(\alpha\beta; q, \omega) \chi_2''(\gamma\delta; q, \omega) |v_{12}(\alpha\beta, \gamma\delta; q, \omega)|^2}{\sinh^2(\hbar\omega/2k_B T)} \quad (5)$$

where χ_i'' is the imaginary part of the random phase approximation irreducible polarizability

$$\chi_i''(\alpha\beta; q, \omega) = \int \frac{d\mathbf{k}}{2\pi} [f_{0\alpha}(\mathbf{k}) - f_{0\beta}(\mathbf{k} + \mathbf{q})] \delta(\varepsilon_\alpha(\mathbf{k}) - \varepsilon_\beta(\mathbf{k} + \mathbf{q}) - \hbar\omega) \quad (6)$$

We shall assume that this form is valid for the remainder of the paper.

B. Interlayer phonon mediated interaction

We assume that the properties of the phonon are the same in both the well and the barrier region. We denote the (bulk) electron-phonon interaction matrix element $M_\lambda(\mathbf{q}, Q_z)$, where λ is the polarization (for clarity the λ will henceforth be suppressed),

Let $\psi_{i\alpha}(z)$ be the transverse wavefunctions of the electrons in well i and subband α . A generalization of the Ref. 11 and 15 to multiple subbands for an effective bare phonon propagator which scatters electrons from $\alpha \rightarrow \beta$ in well i and $\gamma \rightarrow \delta$ in well j gives

$$d_{ij}^{(0)}(\alpha\beta, \gamma\delta; \mathbf{q}, \omega) = \int_{-\infty}^\infty \frac{dQ_z}{2\pi} D^{(0)}(\mathbf{q}, Q_z, \omega) |M(\mathbf{q}, q_z)|^2 f_i(\alpha\beta, Q_z) f_j(\gamma\delta, -Q_z), \quad (7)$$

where $f_i(\alpha\beta, Q_z)$ is the form factor given by

$$f_i(\alpha\beta, Q_z) = \int_{-\infty}^\infty dz \psi_{i\alpha}^*(z) \psi_{i\beta}(z) e^{-iQ_z z} \quad (8)$$

and $D^{(0)}(\mathbf{Q}, \omega)$ is the bare (retarded) bulk phonon Green function. This effective phonon propagator mediates the interaction between the electrons in the adjacent wells.

C. Longitudinal polar optical phonons

The bulk longitudinal polar optic phonon–electron interaction matrix element is given by¹⁷

$$|M^2(Q)| = V_c(Q) \frac{\hbar\omega_{\text{LO}}}{2} \left(\frac{1}{\epsilon_\infty} - \frac{1}{\epsilon_0} \right) \quad (9)$$

where

$$V_c(Q) = \frac{4\pi e^2}{\epsilon_\infty Q^2} \quad (10)$$

is the bare Coulomb interaction.

We assume that the optical phonons have a constant frequency ω_{LO} , since the longitudinal optical phonons have a small dispersion at the Brillouin zone center. Also assuming a finite lifetime γ_{LO}^{-1} gives the form for the bare phonon Green function

$$D_{\text{LO}}^{(0)}(\omega) = \frac{1}{\omega - \omega_{\text{LO}} + i\gamma_{\text{LO}}/2} - \frac{1}{\omega + \omega_{\text{LO}} + i\gamma_{\text{LO}}/2}. \quad (11)$$

With these assumptions, the phonon Green function D can be factored out of Eq. (7), and the spatial form of the effective longitudinal polar optical phonon mediated interaction is exactly the same as the Coulomb interaction. Together with Eqs. (9) and (10) this gives

$$d_{ij,\text{LO}}(\alpha\beta; \gamma\delta; \mathbf{q}, \omega) = D_{\text{LO}}^{(0)}(\omega) \frac{\omega_{\text{LO}}}{2} \left(1 - \frac{\epsilon_\infty}{\epsilon_0} \right) V_{ij}^c(\alpha\beta, \gamma\delta; q) \quad (12)$$

where $V_{ij}^c(\alpha\beta, \gamma\delta; q)$ is the Coulomb matrix element scattering between $\alpha \rightarrow \beta$ in well i and $\gamma \rightarrow \delta$ in well j , together with an in-plane change of $\hbar\mathbf{q}$.

D. Effective screened interlayer interaction

The optical phonon contribution to the drag rate is obtained to lowest order by substituting a $d_{12,\text{LO}}^{(0)}$ into v_{12} in Eq. (2).¹⁸ It has been shown, however, screening of the interaction gives many novel effects in drag experiments.^{14,12,19} In principle screening within the random phase approximation can be included for systems where two subbands participate in each

layer, but one needs to invert an 16×16 matrix to find the screened interaction. However, we shall see that screening is not an important consideration in the case of optical phonon mediated drag.

One important effect of screening is to remove a spurious mathematical divergence in the transresistivity in Eq. (2). If the phonon lifetime is infinite ($\gamma = 0$) then in the absence of screening the bare phonon interaction gives a nonintegrable divergent momentum transfer at the phonon frequency $\omega_{\text{ph}}(q)$ for all q . Even if the lifetime is finite but large, the screening of the electron-phonon interaction is the dominant physical mechanism for elimination of the divergence in the integrand around $\omega_{\text{ph}}(q)$.¹⁵ In the case of the acoustic phonons, the lifetime is very large (on the order of a microsecond) because acoustic phonons are unable to decay via anharmonic terms to other phonons, and hence proper treatment of the screening is important to obtain a quantitatively correct result. On the other hand, the optical phonons can decay via anharmonic terms, leading to much shorter intrinsic lifetimes (on the order of picoseconds). In this case, the broadening of the phonon Green function due to the lifetime is the physically dominant mechanism for the removal of the divergence around $\omega_{\text{ph}}(q)$.

Furthermore, as shown below, most of contribution to the integral Eq. (2) comes from the small q and $\omega \approx \omega_{\text{LO}}$ regime, where the effects of screening are small. Therefore, given the short lifetime of the longitudinal polar optical phonons and the weakness of the screening in the $\omega - q$ space important for optical phonons, it is justified to use the bare interaction $d^{(0)}$ as the interlayer interaction.²⁰

III. CALCULATION AND RESULTS

Longitudinal optical phonons are well defined excitations when γ_{LO} is much smaller than ω_{LO} . Since γ_{LO} is also much smaller than all other frequency scales in the problem, one can approximate

$$|D_{\text{LO}}^{(0)}(\omega)|^2 \approx \frac{2\pi}{\gamma_{\text{LO}}} [\delta(\omega - \omega_{\text{LO}}) + \delta(\omega + \omega_{\text{LO}})]. \quad (13)$$

Note that the real (imaginary) parts of $D_{\text{LO}}^{(0)}(\omega)$ corresponds to the virtual (real) phonon contribution.¹⁵ In the case of optical phonon mediated interaction, the virtual and real phonons contribute equally to the drag.

Using Eq. (13) and Eq. (12) in Eq. (5) and the other approximations stated above, the polar optical phonon mediated drag transresistivity is given by

$$\rho_{21} \approx \frac{1}{8} \frac{\hbar}{e^2} \left(1 - \frac{\epsilon_\infty}{\epsilon_0}\right)^2 \frac{\omega_{\text{LO}}}{\gamma_{\text{LO}}} \frac{\hbar\omega_{\text{LO}}}{k_B T} \frac{1}{\sinh^2(\hbar\omega_{\text{LO}}/2k_B T)} \int_0^\infty d\tilde{q} \tilde{q}^3 \sum_{\alpha\beta\gamma\delta} \chi_1''(\alpha\beta; \tilde{q}, \omega_{\text{LO}}) \chi_2''(\gamma\delta; \tilde{q}, \omega_{\text{LO}}) |V_{12}^c(\alpha\beta, \gamma\delta; \tilde{q})|^2 \quad (14)$$

where $\tilde{q} = q/k_F$. Note that the apparent divergence of the above result $\gamma_{\text{LO}} \rightarrow 0$ is an artifact of the neglect of the dynamical screening scheme, as discussed in the previous section.

We now discuss drag in the case when one has single and double subbands in each layer.

A. Drag in single-subband systems

In the case of drag when there is only a single subband occupied, the expression Eq. (5) reduces to the well-known one given previously.^{9,21} The main difference between previous treatments and optical phonon mediated drag is that optical phonon scattering involves a relatively large exchange of energy, which necessarily implies that the in-plane transfer of momentum is quite large.

In the case when an electron emits or absorbs an optical phonon, its kinetic energy in the direction of the plane changes by $\hbar(\omega_{\text{LO}} - \omega_{\alpha\beta})$, where $\omega_{\alpha\beta}$ is the energy change between the subbands α and β . The momentum transfer $\delta\mathbf{q}$ associated with this for a parabolic band is

$$\delta\mathbf{q} \cdot \langle \mathbf{v} \rangle = \omega_{\text{LO}} - \omega_{\alpha\beta}, \quad (15)$$

where $\langle \mathbf{v} \rangle$ is the velocity of the average of the initial and final state. In the case where the electron stays within the same subband, $\omega_{\alpha\beta} = 0$, and hence $q \gtrsim \omega_{\text{LO}}/\langle v \rangle$.

We insert numbers for GaAs. For density $n = 1.5 \times 10^{11} \text{ cm}^{-2}$, $\langle v \rangle \approx v_F \approx 1.7 \times 10^7 \text{ cm/s}$, and $\omega_{\text{LO}} \approx 6 \times 10^{13} \text{ s}^{-1}$, one obtains a typical q of approximately $3 \times 10^6 \text{ cm}^{-1}$. Given

that the typical distance separating the wells d is on the order of $400 \text{ \AA}$, giving a product $qd \approx 14$. Since the Coulomb interaction (and hence by Eq. (12) the optical phonon mediated interaction) goes as $\exp(-qd)$, and the scattering strength is given by the matrix element squared, one obtains a factor of e^{-28} , which makes it almost impossible to see the effect.

In order to see an effect in single subband systems it is clear that one must reduce the typical momentum transfer $\delta\mathbf{q}$. To do this, one must have a lower optical phonon frequency and a larger $\langle v \rangle$; i.e., a smaller effective mass. A possible candidate is InSb. It has an effective mass of approximately $0.02m_e$ where m_e is the bare electron mass, and a longitudinal optic phonon frequency of 24 meV (compared to $0.067m_e$ and 36 meV , respectively, for GaAs). Assuming an electron density $n = 1.5 \times 10^{11} \text{ cm}^{-2}$, $d = 300 \text{ \AA}$ and well-width $L = 200 \text{ \AA}$, the integral in Eq. (14) for $k_B T \lesssim \hbar\omega_{\text{LO}}$ is approximately 1.9×10^{-3} . The coupling constant $(1 - \epsilon_\infty/\epsilon_0)^2 = 5 \times 10^{-3}$, yielding

$$\rho_{21} \approx 30 \text{ m}\Omega \frac{\hbar\omega_{\text{LO}}}{k_B T} \frac{1}{\sinh^2(\hbar\omega_{\text{LO}}/2k_B T)} \frac{\omega_{\text{LO}}}{\gamma_{\text{LO}}} \quad (16)$$

For $k_B T = \hbar\omega_{\text{LO}}/2$, the temperature terms are equal to 1.44. Therefore, we have for the parameters mentioned above

$$\rho_{\text{InAs}}(T = 110 \text{ K}) \approx 40 \text{ m}\Omega \frac{\omega_{\text{LO}}}{\gamma_{\text{LO}}} \quad (17)$$

The result depends on the product of the optical phonon frequency and lifetime. In a relatively clean system, this ratio is large. Using GaAs as a guide, the lifetime $\tau \approx 7 \text{ ps}$, at 77 K .²² Therefore, $\gamma_{\text{LO}} = 6.6 \times 10^{-16} \text{ eV} / 5 \times 10^{-12} \approx 0.1 \text{ meV}$, giving $\omega_{\text{LO}}/\gamma_{\text{LO}} \approx 360$. This leaves one with a transresistivity of on the order of $1 - 10 \Omega$, which should be experimentally observable.

B. Double subband systems

We define the lower subband as A and upper subband as B . In double-subband systems, one can tune ω_{BA} so that it is equal to ω_{LO} . Then, from Eq. (15) the minimum momentum

transfer is zero, and therefore there can be significant contribution from small in-plane momentum transfer events, without the cut-off in the exponential in $V_{21}^c(q)$. As a consequence, we shall see that the drag rate can be rather large.

We assume the wells are made of GaAs. As mentioned above, the intrasubband excitations do not give significant contributions. Therefore the main contributions come from the terms $\chi_i''(AB; q, \omega)$. Since the integral in Eq. (5) is restricted to $\omega > 0$, the only term is the sum over subband indices which contributes significantly for $k_B T \lesssim \hbar\omega_{LO}$ is $\chi_1(AB; q, \omega)\chi_2(AB; q, \omega)$. In this case, using a square well potential we have

$$V_{12}^c(AB, AB; q) = \left[\frac{16 \cosh(qL/2) \pi^2 qL}{9\pi^2 + 10\pi^2(qL)^2 + (qL)^4} \right]^2 \quad (18)$$

Using the above in Eq. (14) allows us to calculate ρ_{21} . For GaAs, the term $\hbar/(8e^2)(1 - \epsilon_\infty/\epsilon_0)^2 \approx 65 \Omega$. For temperatures $k_B T \lesssim \hbar\omega_{LO}$ at typical experimental densities, the term in the integrand in Eq. (14) is not a strong function of temperature and hence can be evaluated in the $T = 0$ limit (when $k_B T \gtrsim \hbar\omega_{LO}$ the lowest subband becomes nondegenerate and a more careful evaluation of the temperature dependence of χ_i'' in Eq. (14) is necessary). Fig. 2 shows this integral in Eq. (14) as a function of the energy difference ω_{AB} . The drag rate peaks as $\omega_{AB} = \omega_{LO}$ for the reasons stated above. Estimating the ratio $\omega_{LO}/\gamma_{LO} \approx 200$,²² we see that at $T = 100$ K, the optical phonon mediated drag is on the order of 30Ω . As the subband separation moves further away from ω_{LO} , the transresistivity decreases rapidly. The optical phonon mediated interaction is unique in that it is particularly extremely sensitive to width of the wells.

IV. CONCLUSIONS

We have calculated the transresistivity for coupled quantum wells due the mediation of polar optical phonons. The spatial form of the polar optical phonon is identical to that of the Coulomb interaction. We find that in GaAs, when only a single subband is important (i.e., $\hbar\omega_{AB} \gg E_F, k_B T$), the transresistivity is negligible at present experimental

conditions. This is because the polar optical phonon scattering is an inelastic event resulting in a minimum q -vector which substantially reduces the effect. However, it may be possible to see single subband drag in other materials with lower ω_{LO} and lower effective masses such as InSb. In the case where there are two different subbands, the subband energy difference can be engineered to be the same as the ω_{LO} , in which case the optical phonon mediated transresistivity can be relatively large and observable. The magnitude of the drag should be sensitive to the subband energy separation relative to the optical phonon frequencies. It may be possible to observe this resonance by tuning the subband energy splitting through ω_{LO} by means of an external gate which changes the shape of the confining potential in the wells.

In this work we have assumed that the intralayer electron–electron interactions are strong enough that the effective τ is constant (i.e., a drifted Fermi Dirac distribution for all subbands). For the clean samples used in these experiments, this is generally the case. We have also assumed that the phonon frequencies in the barrier and in the well are identical. In fact the optical phonons in $\text{Al}_x\text{Ga}_{1-x}\text{As}$ have frequencies which are, depending of the x , somewhat higher than GaAs. The peak shown in Fig. 2 would tend to be smeared by this mismatch in frequencies.

The author thanks M. C. Bønsager, K. Flensberg, T. J. Gramila, A.-P. Jauho and A. H. MacDonald for useful discussions and comments.

REFERENCES

- ¹ M. B. Pogrebinskii, Fiz. Tekh. Poluprovodn. **11**, 637 (1977) [Sov. Phys. Semicond. **11**, 372 (1977)].
- ² P. J. Price, Physica B **117**, 750 (1983).
- ³ T. J. Gramila, J. P. Eisenstein, A. H. MacDonald, L. N. Pfeiffer, and K. W. West, Phys. Rev. Lett. **66**, 1216 (1991); Surf. Sci. **263**, 446 (1992); Phys. Rev. B **47**, 12957 (1993); Physica B **197**, 442 (1994).
- ⁴ U. Sivan, P. M. Solomon, and H. Shtrikman, Phys. Rev. Lett. **68**, 1196 (1992).
- ⁵ H. Rubel, E. H. Lindfield, D. A. Ritchie, K. M. Brown, M. Pepper and G. A. C. Jones, Semicond. Sci. Technol. **10**, 1229 (1995); Surf. Sci. **361/362** 134 (1996).
- ⁶ H. Rubel, A. Fisher, W. Ditcher, K. von Klitzing and K. Eberl, Phys. Rev. Lett. **78**, 1763 (1997).
- ⁷ N. P. R. Hill, J. T. Nicholls, E. H. Linfield, M. Pepper, D. A. Ritchie, A. R. Hamilton and G. A. C. Jones, J. Phys. Cond. Mat. **8**, L557 (1996).
- ⁸ N. P. R. Hill, J. T. Nicholls, E. H. Linfield, M. Pepper, D. A. Ritchie, G. A. C. Jones, B. Y.-K. Hu and K. Flensberg, Phys. Rev. Lett. **78**, 2204 (1997).
- ⁹ A.-P. Jauho and H. Smith, Phys. Rev. B **47**, 4420 (1993).
- ¹⁰ H. C. Tso, P. Vasilopoulos, and F. M. Peeters, Phys. Rev. Lett. **68**, 2516 (1992).
- ¹¹ C. Zhang and Y. Takahashi, J. Phys. Cond. Mat. **5**, 5009 (1993).
- ¹² K. Flensberg and B. Y.-K. Hu, Phys. Rev. Lett. **73**, 3572 (1994).
- ¹³ L. Świerkowski, J. Szymanski, and Z. W. Gortel, Phys. Rev. Lett. **74**, 3245 (1995).
- ¹⁴ K. Flensberg and B. Y.-K. Hu, Phys. Rev. B **52**, 14796 (1995).
- ¹⁵ M. C. Bønsager, K. Flensberg, B. Y.-K. Hu and A. H. MacDonald (unpublished).

- ¹⁶ B. Y.-K. Hu and K. Flensberg, Phys. Rev. B **53**, 10072 (1996).
- ¹⁷ See e.g., G. D. Mahan, Many-Particle Physics 2nd. Ed. (Plenum Press, New York, 1990).
- ¹⁸ In principle, the bare interlayer interaction v_{12} is the sum of the Coulomb term V_{12}^c and phonon term d , and there are interference terms between these two terms. However, the optical phonon contribution is strong and sharp and localized in $q - \omega$ space, and hence the interference effects $V_{12}^c d$ are minimal compared to the “purely phonon” term d^2 .
- ¹⁹ M. C. Bønsager, K. Flensberg, B. Y.-K. Hu, and A.-P. Jauho, Phys. Rev. Lett. **77**, 1366 (1996).
- ²⁰ One possible screening effect which might complicate the situation is coupling the optical phonons to intersubband plasmons (A. Kuznetsov, private communication). As the intersubband plasmons always lie above the intersubband single-particle continuum due to the depolarization shift, they do not directly participate in the drag, but they can couple to the optical phonons and shift the frequency of that mode. The main qualitative effects mentioned in this paper, however, should not be altered.
- ²¹ L. Zheng and A. H. MacDonald, Phys. Rev. B **48**, 8203 (1993).
- ²² B. K. Ridley, J. Phys. Cond. Mat. **8**, 2511 (1996); E. Haro-Poniatowski, J. L. Escamilla-Reyes and K. H. Wanser, Phys. Rev. B **53**, 12121 (1996); A. R. Bhatt, K. W. Kim, and M. A. Stroscio, J. App. Phys. **76**, 3905 (1994).

FIGURES

FIG. 1. Schematic for the double quantum wells system with two subbands, denoted A for lower and B for upper. Experimentally current is passed through layer 1 and a voltage is measured in layer 2. The phonon-mediated scattering is indicated by the Feynman diagram. The filled dots are the electron-phonon vertices and the dotted line indicates the phonon Green function. The $\alpha\beta\gamma\delta$ are the subband indices, $\mathbf{k}, \mathbf{k}'$ are the in-plane momenta before collision and $\mathbf{q}$ is the in-plane momentum transfer.

FIG. 2. The integral in Eq. (14), $d = 400 \text{ \AA}$, $n = 1.5 \times 10^{11} \text{ cm}^{-2}$, and $\omega_{\text{LO}} = 36 \text{ meV} \approx 400 \text{ K}$, as in GaAs. We assume infinite square wells for the confinement, in which case $\omega_{AB} = \omega_{\text{LO}}$ for $L \approx 215 \text{ \AA}$. Inset: temperature dependence of the transresistivity for low temperatures, $x/\sinh^2(1/2x)$ as a function of $x = \hbar\omega_{\text{LO}}/k_B T$ [see Eqs. (14) and (16)].



